

EXACTNESS FROM PROPER ACTIONS

J. BRODZKI, G.A. NIBLO, AND N.J. WRIGHT

ABSTRACT. In this paper we show that if a discrete group G acts properly isometrically on a discrete space X for which the uniform Roe algebra $C_u^*(X)$ is exact then G is an exact group. As a corollary, we note that if the action is cocompact then the following are equivalent: The space X has Yu's property A; $C_u^*(X)$ is exact; $C_u^*(X)$ is nuclear.

INTRODUCTION

It is a remarkable result of Yu that any discrete metric space with bounded geometry that satisfies a Følner-type condition, which he called property A, also satisfies the coarse Baum-Connes conjecture [7]. The case of a countable discrete group, regarded as a coarse metric space of bounded geometry, was studied by Higson and Roe [2], Guentner and Kaminker [1], and Ozawa [3]. They showed that Yu's property A is equivalent to exactness of its reduced C^* -algebra $C_r^*(G)$ and to nuclearity of its uniform Roe algebra $C_u^*(G)$. Furthermore, Roe showed for a discrete bounded geometry metric space X that if X has property A then $C_u^*(X)$ is nuclear ([4], Proposition 11.41). It is tempting to conjecture therefore that for a discrete metric space of bounded geometry, nuclearity of $C_u^*(X)$ is in fact equivalent to property A. This conjecture holds if X admits a free cocompact action by a countable group G : since the action is free we may identify $C_u^*(G)$ with a subalgebra of $C_u^*(X)$. If we assume that $C_u^*(X)$ is nuclear, then $C_u^*(G)$ will be exact and G will have property A. The fact that G acts cocompactly on X implies that X and G are coarsely equivalent so X also has property A.

The conjecture is more delicate than it may appear since while property A is a coarse invariant, the uniform Roe algebra is not. For example, for a finite space X with n points, $C_u^*(X)$ is the algebra of $n \times n$ matrices; however all finite spaces are coarsely equivalent to a point.

In this paper we address the case of the conjecture when the space admits a proper, cocompact isometric action by a countable group G . Generalising a theorem in coarse geometry from the case of a free action to a proper action usually requires only a minor adjustment of the argument. This is not the case in our context. The fact that the uniform Roe algebra is not a coarse invariant is manifested in the observation that when the action is not free we no longer have the required embedding of $C_u^*(G)$ into $C_u^*(X)$. The key idea of this paper is to replace the proper action on X by a partially defined free action on an orbit of the original action. In outline we proceed as follows.

Given an orbit Y of the action of G on X and a splitting ϕ of the natural surjection $G \rightarrow Y$ we identify the algebra $C_u^*(Y)$ with the subalgebra $C_u^*(\phi(Y))$ of $C_u^*(G)$. We construct a free partial action of G on the orbit Y defined in terms of

the right action of G on itself and the identification of $\phi(Y)$ with Y . Finally we may appeal to coarse equivalences to establish the following.

Theorem 1. *Let X be a uniformly discrete metric space, and let G be a countable group acting properly by isometries on X . If $C_u^*(X)$ is an exact C^* -algebra, then G is an exact group.*

From this we deduce the following equivalence.

Corollary 2. *If X is a uniformly discrete metric space admitting a proper, cocompact isometric action by a countable group G then the following are equivalent.*

- (1) X has property A;
- (2) $C_u^*(X)$ is nuclear;
- (3) $C_u^*(X)$ is exact;
- (4) G is exact.

Proof. of Corollary 2

The implication $1 \Rightarrow 2$ was established by Roe in [4, Prop. 11.41]. That 2 implies 3 is a well known result, see, e.g. [6]. Theorem 1 provides the implication $3 \Rightarrow 4$. Finally, as the action of G on X is cocompact, X and G are coarsely equivalent, which gives the implication $4 \Rightarrow 1$. $\square$

BACKGROUND

Throughout the paper we will assume that G is a countable group equipped with the unique (up to coarse equivalence) proper left invariant metric d_G . For $R \geq 0$ we will use the notation $B_R(g)$ to denote the closed R -ball about g in G .

Definition 3. A uniformly discrete metric space (X, d_X) has *property A* if for all $R, \varepsilon > 0$ there exists a family of finite non-empty subsets A_x of $X \times \mathbb{N}$, indexed by x in X , such that

- for all x, y with $d_X(x, y) < R$ we have $\frac{|A_x \Delta A_y|}{|A_x \cap A_y|} < \varepsilon$;
- there exists S such that for all x and $(y, n) \in A_x$ we have $d_X(x, y) \leq S$.

Definition 4. A kernel $u: X \times X \rightarrow \mathbb{R}$ has (R, ε) -*variation* if $d_X(x, y) \leq R$ implies $|u(x, y) - 1| < \varepsilon$.

Theorem 5 ([5], Proposition 3.2). *If X is a bounded geometry discrete metric space, then X has property A if and only if for all $R, \varepsilon > 0$ there exists a positive definite kernel u with R, ε variation, and such that there exists S for which $d_X(x, y) > S$ implies $u(x, y) = 0$.*

Proposition 6. *Let G be a countable group acting properly isometrically on a proper metric space X . Then for any basepoint x_0 in X , the map $\psi: g \mapsto gx_0$ is a uniform embedding, and any map $\phi: Gx_0 \rightarrow G$ such that $\psi \circ \phi = \text{id}_{Gx_0}$ is a coarse inverse to ψ . If moreover G acts cocompactly then ψ is a coarse equivalence between G and X .*

Proof. First we verify that ψ is a coarse map. For g, h in G , we have $d_X(gx_0, hx_0) = d_X(x_0, g^{-1}hx_0)$ as the action is isometric. Given R , if $d_G(g, h) \leq R$ then $g^{-1}h$ lies in $B_R(e)$ which, by properness of the metric d_G is finite, so $d_X(x_0, g^{-1}hx_0)$ is bounded by some number S . Thus for all R there exists S such that $d_G(g, h) \leq R$ implies $d_X(\psi(g), \psi(h)) \leq S$. Properness of ψ follows from properness of the action, so ψ is a coarse map.

Now let $\phi: Gx_0 \rightarrow G$ be a splitting of ψ . Then for x, y in Gx_0 we have $x = \phi(x)x_0, y = \phi(y)x_0$. If $d_X(x, y) \leq R$ then $d_X(x_0, \phi(x)^{-1}\phi(y)x_0) \leq R$. Properness of the action ensures that there are only finitely many elements g of G with $d_X(x_0, gx_0) \leq R$, so $\phi(x)^{-1}\phi(y)$ lies in $B_S(e)$, for some S , i.e. $d_G(\phi(x), \phi(y)) \leq S$. The map ϕ is injective, thus it is proper, so ϕ is also a coarse map.

The composition $\psi \circ \phi$ is the identity on Gx_0 by definition, while $g^{-1}(\phi \circ \psi(g))$ is an element of G fixing x_0 . The stabiliser of x_0 is finite, so $d_G(g, \phi \circ \psi(g))$ is bounded as a function of g , i.e. $\phi \circ \psi$ is close to the identity on G . Thus ψ is a uniform embedding from G to X , and ψ, ϕ implement a coarse equivalence between G and its image.

If the action is cocompact then for some R , the G translations of the R -ball about x_0 cover X , so Gx_0 is R -dense in X . It follows immediately that the inclusion of Gx_0 into X is a coarse equivalence, so $\psi: G \rightarrow X$ is a coarse equivalence. $\square$

Definition 7. A kernel $k: X \times X \rightarrow \mathbb{C}$ has *finite propagation* if there exists $R \geq 0$ such that $k(x, y) = 0$ for $d(x, y) > R$. The *propagation* of k is the smallest such R .

If X is a proper discrete metric space, and k is a finite propagation kernel then for each x there are only finitely many y with $k(x, y) \neq 0$. Thus k defines a linear map from $l^2(X)$ to itself, $k * v(x) = \sum_{y \in X} k(x, y)v(y)$. These linear maps are also said to have finite propagation. Note that if additionally X has bounded geometry, then every bounded finite propagation kernel gives rise to a *bounded* operator on $l^2(X)$.

Definition 8. The *uniform Roe algebra*, $C_u^*(X)$, is the C^* -algebra completion of the algebra of bounded operators on $l^2(X)$ having finite propagation.

Definition 9. A *partial translation* of X is a subset t of $X \times X$ such that the coordinate projections of t onto X are injective, and such that $d_X(x, y)$ is bounded for $(x, y) \in t$.

A partial translation can be viewed as a partially defined map from X to X which is close to the identity (where defined). Therefore a partial translation gives rise to a partial isometry of $l^2(X)$ which has finite propagation and hence is an element of $C_u^*(X)$. By definition, the partial isometry is defined by $t(\delta_y) = \delta_x$ if $t(y) = x$, and $t(\delta_y) = 0$ if $t(y)$ is undefined.

PROOF OF THEOREM 1

Fix a basepoint x_0 in X , and let Y be the orbit of x_0 under the action of G . Given a finite propagation operator on $l^2(Y)$ we can extend it to $l^2(X) = l^2(Y) \oplus l^2(X \setminus Y)$ by defining it to be zero on $l^2(X \setminus Y)$. Thus the algebra $C_u^*(Y)$ is a subalgebra of $C_u^*(X)$, hence exactness for $C_u^*(X)$ implies exactness for $C_u^*(Y)$.

For each $y \in Y$, pick an element $\phi(y)$ of G such that $\phi(y)x_0 = y$. We construct a partially defined action of the group G on the space Y as follows. For $g \in G, y \in Y$ we define $g \diamond y = x$ if and only if $\phi(y)g^{-1} = \phi(x)$. The element $g \diamond y$ in Y is uniquely determined if it exists, since then $g \diamond y = \phi(y)g^{-1}x_0$. However it will be undefined if $\phi(y)g^{-1}$ is not in the image of ϕ . Note that $g \diamond$, viewed as a partially defined map from X to X is a partial translation, since

$$d_G(\phi(y), \phi(g \diamond y)) = d_G(\phi(y), \phi(y)g^{-1}) = d_G(e, g^{-1})$$

which is independent of y , and ϕ is a coarse equivalence by Proposition 6. Note that in the case where G acts *freely*, the map ϕ is uniquely determined and is a bijection between Y and G . Using this to identify Y with G , the action $g\Diamond$ is identified with the action of G on itself by right multiplication by g^{-1} .

Given R, ε we will produce a positive kernel u on Y with (R, ε) -variation, and satisfying the hypothesis that $u(x, y)$ vanishes for x, y far apart. Let E_R be the set of elements of G of the form $\phi(x)^{-1}\phi(y)$, x, y in Y with $d_X(x, y) \leq R$. As ϕ is a coarse map, there exists S such that if $d_X(x, y) \leq R$ then $d_G(\phi(x), \phi(y)) \leq S$, so E_R is contained in the ball $B_S(e)$ in G , in particular it is finite. Elements of G act as partial translations on Y via $\Diamond$, hence we can identify E_R with a finite subset of $C_u^*(Y)$.

Using the characterisation of exactness ([3] Lemma 2), there exists a completely positive finite rank map $\theta: C_u^*(Y) \rightarrow B(l^2(Y))$ such that:

- (1) θ has the form $\theta(\cdot) = \sum_{i=1}^d \langle \delta_{a_i}, \cdot \delta_{b_i} \rangle T_i$, for some a_i, b_i in Y , and T_i in $B(l^2(Y))$, and
- (2) for all g in E_R we have $\|\theta(g) - g\| < \varepsilon$.

Let F_R denote the set $\{a_i, b_i : i = 1, \dots, d\}$, and note that for any partial translation t such that the image of F_R under t does not meet F_R , we have $\theta(t) = 0$. We now define

$$u(x, y) = \langle \delta_x, \theta(\phi(x)^{-1}\phi(y))\delta_y \rangle \text{ for } x, y \text{ in } Y,$$

where the elements $\phi(x)^{-1}\phi(y)$ of G are regarded as elements of $C_u^*(Y)$ as above.

First we will verify positivity of u ; this is not immediate because as an operator on $l^2(X)$, $\phi(x)^{-1}\phi(y)$ is not necessarily the composition of the operators corresponding to $\phi(x)^{-1}$ and $\phi(y)$. This is because $(\phi(x)^{-1}\phi(y)) \Diamond y'$ may be defined even when $\phi(x)^{-1} \Diamond (\phi(y) \Diamond y')$ is not.

For each y in Y , we define a bounded linear map s_y from $l^2(Y)$ to $l^2(G)$ as follows. Let $s_y(\delta_{y'}) = \delta_g$ where $g = \phi(y')\phi(y)^{-1}$ in G . Then its adjoint s_y^* satisfies $s_y^*(\delta_g) = \delta_{x'}$ if there exists x' with $\phi(x')\phi(x)^{-1} = g$, (such an x' must be unique by injectivity of ϕ) and is zero otherwise. Thus for x, y, y' in Y , the vector $s_x^*s_y(\delta_{y'})$ is $\delta_{x'}$ for some $x' \in Y$ if we have

$$\phi(y')\phi(y)^{-1} = \phi(x')\phi(x)^{-1},$$

and it is zero otherwise. Note that we can rewrite this as $\phi(y')(\phi(x)^{-1}\phi(y))^{-1} = \phi(x')$ i.e. $x' = (\phi(x)^{-1}\phi(y)) \Diamond y'$. We conclude that

$$\begin{aligned} s_x^*s_y(\delta_{y'}) &= \delta_{x'} = (\phi(x)^{-1}\phi(y))(\delta_{y'}) & \text{if } x' = (\phi(x)^{-1}\phi(y)) \Diamond y', \\ s_x^*s_y(\delta_{y'}) &= 0 = (\phi(x)^{-1}\phi(y))(\delta_{y'}) & \text{if } (\phi(x)^{-1}\phi(y)) \Diamond y' \text{ is undefined.} \end{aligned}$$

Hence $s_x^*s_y = \phi(x)^{-1}\phi(y)$ as an operator on $l^2(Y)$. The operators $(\phi(x)^{-1}\phi(y)) = (s_x^*s_y)$ therefore form a positive matrix over Y , so positivity of u follows from complete positivity of θ .

We will now show that u has (R, ε) -variation. For x, y with $d_X(x, y) \leq R$ we have $\phi(x)^{-1}\phi(y)$ in E_R , hence it follows from (2) above that

$$\|\phi(x)^{-1}\phi(y) - \theta(\phi(x)^{-1}\phi(y))\| < \varepsilon.$$

We recall that $(\phi(x)^{-1}\phi(y)) \Diamond y'$ is defined if $\phi(y')(\phi(x)^{-1}\phi(y))^{-1}$ is in the image of ϕ . In particular for $y' = y$ we have that $\phi(y)(\phi(x)^{-1}\phi(y))^{-1} = \phi(x)$ which is in the

image of ϕ , so $(\phi(x)^{-1}\phi(y)) \diamond y$ is defined and equals x . Thus $\phi(x)^{-1}\phi(y)(\delta_y) = \delta_x$, so $\langle \delta_x, \phi(x)^{-1}\phi(y)(\delta_y) \rangle = 1$. Hence

$$|1 - u(x, y)| = |\langle \delta_x, (\phi(x)^{-1}\phi(y) - \theta(\phi(x)^{-1}\phi(y)))(\delta_y) \rangle| < \varepsilon.$$

We conclude that u has (R, ε) -variation as required.

Finally we will show that $u(x, y)$ vanishes for $d_X(x, y)$ sufficiently large. As ϕ is a uniform embedding, for all R' there exists S' such that if $d_G(\phi(x), \phi(y)) \leq R'$ then $d_X(x, y) \leq S'$. As F_R is a finite subset of Y we note that $\{\phi(x')^{-1}\phi(y') : x', y' \in F_R\}$ is a finite subset of G , and choose R' such that this is contained in the ball $B_{R'}(e)$ in G . Now if $x, y \in Y$ and $x', y' \in F_R$ with $(\phi(x)^{-1}\phi(y)) \diamond y' = x'$ then $\phi(y')(\phi(x)^{-1}\phi(y))^{-1} = \phi(x')$ so $\phi(x')^{-1}\phi(y') = \phi(x)^{-1}\phi(y)$. Hence

$$d_G(e, \phi(x)^{-1}\phi(y)) = d_G(e, \phi(x')^{-1}\phi(y')) \leq R',$$

i.e. $d_G(\phi(x), \phi(y)) \leq R'$, so $d_X(x, y) \leq S'$. Thus if $d_X(x, y) > S'$ then for y' in F_R , $(\phi(x)^{-1}\phi(y)) \diamond y'$ cannot be an element of F_R . It follows that if $d_X(x, y) > S'$ then $\langle \delta_{x'}, (\phi(x)^{-1}\phi(y))\delta_{y'} \rangle = 0$ for all x', y' in F_R , so $\theta(\phi(x)^{-1}\phi(y))$ vanishes. Hence $u(x, y)$ also vanishes as required.

We have shown that for all R, ε there exists a positive kernel on Y with (R, ε) -variation, and which vanishes for x, y far apart. It follows by Theorem 5 that Y has property A. As G is coarsely equivalent to Y , it too has property A, so by Ozawa's result G is exact.

REFERENCES

- [1] E. Guentner and J. Kaminker, *Exactness and the Novikov conjecture*, Topology 41 (2002), no. 2, 411–418.
- [2] N. Higson and J. Roe, *Amenable group actions and the Novikov conjecture*, J. Reine Angew. Math. 519 (2000), 143–153.
- [3] N. Ozawa, *Amenable actions and exactness for discrete groups*, C. R. Acad. Sci. Paris Sér. I Math. 330 (2000), no. 8, 691–695.
- [4] J. Roe, *Lectures on coarse geometry* University Lecture Series 31, American Mathematical Society (2003).
- [5] J.-L. Tu, *Remarks on Yu's "property A" for discrete metric spaces and groups*, Bull. Soc. Math. France 129 (2001), 115–139.
- [6] S. Wassermann, *Exact C^* -algebras and related topics*, Seoul National University, Seoul 1994.
- [7] G. L. Yu, *The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into a Hilbert space*, Inventiones Mathematicae 138 (2000), 201–240.

E-mail address: j.brodzki@soton.ac.uk

E-mail address: g.a.niblo@soton.ac.uk

E-mail address: n.j.wright@soton.ac.uk

SCHOOL OF MATHEMATICS, UNIVERSITY OF SOUTHAMPTON, HIGHFIELD, SO17 1BJ